

Published in final edited form as:

Biochemistry. 2010 March 30; 49(12): 2657–2671. doi:10.1021/bi901921k.

P90 RIBOSOMAL S6 KINASE 2, A NOVEL GPCR KINASE, IS REQUIRED FOR GROWTH FACTOR-MEDIATED ATTENUATION OF GPCR SIGNALING†

Ryan T. Strachan[‡], John A. Allen[§], Douglas J. Sheffler^{||}, and Bryan L. Roth^{§,*}

[‡]Department of Biochemistry, Case Western Reserve University Medical School Cleveland, OH 44106

[§]Department of Pharmacology, University of North Carolina at Chapel Hill Medical School, Chapel Hill, NC 27599

^{||}Department of Pharmacology, Vanderbilt University, Nashville, TN 37232.

Abstract

The 5-hydroxytryptamine 2A (5-HT_{2A}) receptor is a member of the G protein-coupled receptor superfamily (GPCR) and plays a key role in transducing a variety of cellular signals elicited by serotonin (5-HT; 5-hydroxytryptamine) in both peripheral and central tissues. Recently, we discovered that the ERK/MAPK effector p90 ribosomal S6 kinase 2 (RSK2) phosphorylates the 5-HT_{2A} receptor and attenuates 5-HT_{2A} receptor signaling. This raised the intriguing possibility of a regulatory paradigm whereby receptor tyrosine kinases (RTKs) attenuate GPCR signaling (i.e., ‘inhibitory cross-talk’) by activating RSK2 [Strachan et al. (2009) *J. Biol. Chem.* 284, 5557-5573]. We report here that activation of multiple endogenous RTKs such as the epidermal growth factor receptor (EGFR), the platelet-derived growth factor receptor (PDGFR), and ErbB4 significantly attenuates 5-HT_{2A} receptor signaling in a variety of cell types including mouse embryonic fibroblasts (MEFs), mouse vascular smooth muscle cells (mVSMCs), and primary cortical neurons. Importantly, genetic deletion of RSK2 completely prevented signal attenuation, thereby suggesting that RSK2 is a critical mediator of inhibitory cross-talk between RTKs and 5-HT_{2A} receptors. We also discovered that P2Y purinergic receptor signaling was similarly attenuated following EGFR activation. By directly testing multiple endogenous growth factors/RTK pathways and multiple Gq-coupled GPCRs, we have now established a cellular mechanism whereby RTK signaling cascades act via RSK2 to attenuate GPCR signaling. Given the pervasiveness of growth factor signaling, this novel regulatory mechanism has the potential to explain how 5-HT_{2A} receptors are regulated in vivo, with potential implications for human diseases in which 5-HT_{2A} or RTK activity is altered (e.g. neuropsychiatric and neurodevelopmental disorders).

The GPCR superfamily mediates essential functions in organisms as diverse as unicellular choanoflagellates and humans (1, 2). In humans, GPCRs comprise approximately 2% of the genome to transduce signals elicited by both endogenous and exogenous ligands (3-5). Not

[†]This work was supported by RO1MH61887, NO1MH32004, U19MH82441, T32HD040127, and UNC Neurodevelopmental Disorders Research Center.

^{*}**Corresponding Author Footnote.** Department of Pharmacology 4072 Genetics Medicine Building CB 7365 Chapel Hill, NC 27599-7365. Phone: 919-966-7535; Fax: 919-843-5788; bryan_roth@med.unc.edu . ryan.strachan@receptor-biol.duke.edu

Supporting Information. Supporting information is available showing (1) microarray analysis of genes involved in IR and IGF-1 R signaling pathways in RSK2+/+ and RSK2-/- MEFs and (2) IGF-1 activation of the IGF-1 R. This material is available free of charge via the internet at <http://pubs.acs.org>.

surprisingly, GPCR dysregulation is associated with many human diseases (6), thus explaining why GPCRs are successful therapeutic targets and remain the focus of intense drug discovery efforts (7).

The Gq-coupled 5-HT_{2A} receptor, in particular, is one of 14 GPCRs that mediates the pleiotropic actions of 5-HT in both peripheral and central tissues (8, 9). The 5-HT_{2A} receptor is an important therapeutic target for a large number of psychiatric and medical diseases (9), and is also the site of action of most, but not all hallucinogens which function as 5-HT_{2A} receptor agonists (10, 11)(Keiser et al., 2009, in press). Additionally, atypical antipsychotics (e.g., clozapine) are thought to mediate their therapeutic actions, at least in part, by antagonizing 5-HT_{2A} receptors (12).

We recently discovered a novel regulatory mechanism whereby RSK2 interacts with 5-HT_{2A} serotonin receptors and attenuates receptor signaling *via* direct receptor phosphorylation (13, 14). RSK2 is a multifunctional ERK/MAPK effector activated downstream of growth factor signal cascades involving RTKs (15). This raised the intriguing possibility of a new regulatory mechanism whereby RTKs attenuate GPCR signaling (referred to here as ‘inhibitory cross-talk’) by activating RSK2. These studies led to the initial discovery that activation of the EGFR attenuates 5-HT_{2A} receptor signaling, presumably *via* RSK2 activation (14). These preliminary data were intriguing for several reasons including: (1) they suggested for the first time that the 5-HT_{2A} receptor is part of an emerging regulatory paradigm whereby activated RTKs attenuate GPCR signaling (16-22), and (2) they were the first to identify RSK2 as a novel mediator of inhibitory cross-talk between growth factor-activated RTKs and a GPCR.

In this paper, we show that activation of various endogenous RTKs (i.e. EGFR, PDGFR, and ErbB4) significantly attenuates 5-HT_{2A} receptor signaling in multiple cell types (i.e., in MEFs, mVSMCs, and primary cortical neurons). In contrast, insulin-like growth factor 1 (IGF-1), which only weakly activates RSK2, fails to attenuate 5-HT_{2A} receptor signaling. Together with evidence that genetic deletion of RSK2 is sufficient to prevent RTK-mediated signal attenuation in all tested cellular backgrounds, these findings support a novel role for RSK2 in inhibitory cross-talk between RTKs and the 5-HT_{2A} receptor. Significantly, we also discovered that P2Y purinergic receptor signaling, which is regulated by RSK2, was similarly attenuated following EGF receptor activation in wild-type (RSK2+/+) MEFs. By testing several endogenous growth factors/RTK pathways and multiple Gq-coupled GPCRs, we have now established a cellular mechanism whereby RTK signaling cascades attenuate GPCR signaling through RSK2. These findings provide an initial framework for a conserved regulatory mechanism whereby RTKs act *via* RSK2 to attenuate GPCR signaling, and given the complexity of cellular signaling, have the potential to explain how these receptors are regulated *in vivo*.

Moreover, because null mutations of RSK2 lead to Coffin-Lowry Syndrome which exhibits behaviors characteristic of 5-HT_{2A} dysregulation including a schizophrenia-like psychosis and cognitive impairment (23), these findings may explain, in part, some of the clinical manifestations of this neurodevelopmental disease..

Experimental Procedures

Materials

Cell culture reagents including fetal bovine serum (FBS), Dulbecco’s modified essential medium (DMEM), Trypsin-EDTA, F-12 nutrient mixture, OptiMEM, neurobasal medium, B27 supplement, Hank’s balanced salt solution (HBSS), sodium pyruvate, penicillin, and streptomycin were supplied by Gibco (Invitrogen, Carlsbad, CA). Serotonin, 5-methoxy-

N,N-dimethyltryptamine (5-methoxyDMT), human EGF, human PDGF-AB and PDGF-BB, human TGF- α , IGF-1, papain, probenecid, bovine serum albumin (BSA), low molecular weight poly-L-lysine, sodium tetraborate, L-glutamine, and all other standard reagents were supplied by Sigma-Aldrich Corp. (St. Louis, MO). Boric acid was supplied by EMD Chemicals (Gibbstown, NJ). MDL100907 and lisuride were acquired as previously detailed (12). Collagenase II was obtained from Worthington Biochemical Corp. (Lakewood, NJ) and elastase (grade II) was supplied by Roche Applied Science (Indianapolis, IN). Restriction endonucleases were supplied by New England Biolabs (Ipswich, MA). [^3H]-Ketanserin was obtained from PerkinElmer Life and Analytical Sciences (Waltham, MA). Protein A/G agarose was supplied by Santa Cruz Biotechnology, Inc. (Santa Cruz, CA).

Serum Dialysis

We extensively dialyzed FBS to remove the 5-HT present in serum. Briefly, 500 mL FBS was placed into dialysis tubing (Spectra/Por 3500 MWCO, Spectrum Laboratories, Rancho Dominguez, CA) and equilibrated with 4 L of cold dialysis buffer (120 mM NaCl, 10 mM Tris-HCl, pH 7.5 at RT) for 24 hr at 4°C with stirring. The buffer was changed five times totaling 120 hr of dialysis. The dialyzed FBS was then sterile-filtered (0.22 μm , Millipore) and aliquots were stored at -20°C until further use. HPLC-electrochemical detection analysis of the dialyzed serum determined that, when used at a concentration of 5%, our dialyzed culture medium contained 33 times less 5-HT than commercially dialyzed FBS (i.e., 0.039 nM vs. 1.3nM 5-HT).

Cell Culture and Transfection

The RSK2 $^{+/+}$ and RSK2 knockout (RSK2 $^{-/-}$) MEFs stably expressing similar levels of 5-HT $_2\text{A}$ receptors were generated previously by Sheffler et al. (13) using MEFs originally isolated from RSK2 $^{+/+}$ and RSK2 $^{-/-}$ mice (24). Mouse VSMCs and cortical neurons were isolated as detailed below. HEK293T cells were obtained from the American Type Culture Collection (Manassas, VA). All cell lines were cultured at 37°C in a humidified environment in the presence of 5% CO $_2$. Specifically, HEK293T and mVSMC cell lines were maintained in standard medium (DMEM supplemented with 10% FBS, 1 mM sodium pyruvate, 100 units/mL penicillin, and 100 $\mu\text{g}/\text{mL}$ streptomycin). Polyclonal populations of MEFs stably expressing FLAG-tagged rat 5-HT $_2\text{A}$ receptors were cultured in standard medium supplemented with 4 $\mu\text{g}/\text{mL}$ puromycin to maintain selection pressure. Primary cortical neurons were maintained in complete neurobasal medium (neurobasal medium, 1 \times B27 supplement, 0.5 mM L-glutamine, 25 μM glutamate, 100 units/mL penicillin, and 100 $\mu\text{g}/\text{mL}$ streptomycin). Eugene6 (Roche) was used exactly as described by the manufacturer to transiently transfect sub-confluent HEK 293T cells.

cDNA Constructs

For generation of stable cell lines, the rat 5-HT $_2\text{A}$ receptor containing a cleavable N-terminal *H. influenza* hemagglutinin membrane insertion signal sequence (25) and N-terminal FLAG (DYKDDDDK) affinity tag (FLAG-5-HT $_2\text{A}$) (26) was subcloned into the pBABE retroviral vector containing a puromycin resistance gene (FLAG-5-HT $_2\text{A}$ -pBABEpuro) (27). Briefly, 5' EcoRI and 3' SalI restriction sites were introduced into the FLAG-5-HT $_2\text{A}$ sequence via the following PCR primers: 5'AAAGAATTCGCCACCATGAAGACGATCAT3' (*EcoRI* highlighted) and 5'AAAGTCGACTCACACACAGCTAACCTTTTC3' (*SalI* highlighted). The FLAG-5-HT $_2\text{A}$ -pBABEpuro construct was sequence-verified (Case Western Reserve University Genomics Core Facility, Cleveland, OH) and determined via competition radioligand binding assays to bind 5-HT with characteristic affinity (<http://pdsp.med.unc.edu/pdsp.php>)(28).

For infection of primary cortical neurons, the rat 5-HT_{2A} receptor containing the green fluorescent protein (GFP) inserted between amino acids 452 and 453 within the C-terminus (5-HT_{2A}-GFP-CT) (29) was subcloned into the FUGW lentiviral construct (5-HT_{2A}-GFP-CT-FUGW) (30). Briefly, 5' XbaI and 3' EcoRI restriction sites were introduced into the 5-HT_{2A}-GFP-CT sequence via the following primers:

5'AAAATCTAGAGCCACCATGGAAATTCTTTGTGAAG3' (*XbaI* highlighted) and 5'TTTTGAATTCTCACACACAGCTAACCTTTTCATTC3' (*EcoRI* highlighted). The resulting 5-HT_{2A}-GFP-CT-FUGW construct was sequence-verified by automated sequencing (UNC-Chapel Hill DNA sequencing facility, Chapel Hill, NC).

Microarray Analysis and Pathway Generation

Microarray studies were performed previously by Sheffler et al. (13) to compare gene expression profiles in RSK2^{+/+} and RSK2^{-/-} MEFs. For pathway analysis of RSK2^{-/-} and RSK2^{+/+} fibroblast gene expression patterns, GenMAPP and MAPPFinder software packages were used as previously detailed (13, 31, 32).

Isolation of Mouse Aortic Vascular Smooth Muscle Cells

Mouse aortic VSMCs were isolated from 12-week old mice (three mice per genotype), as previously detailed (33). Briefly, mice were sacrificed by cervical dislocation and immediately perfused with 25 mL of 1× HBSS (without Ca²⁺ and Mg²⁺). Under sterile conditions, the abdominal/thoracic aorta extending from the ilial bifurcation to aortic arch was carefully microdissected and rinsed with HBSS. The pooled aorta were then incubated with collagenase buffer (175 U/mL in HBSS, filtered through 0.2 µm polyethersulfone membrane) for 15 min at 37°C in the presence of 5% CO₂. After the adventitial layer was removed, the aorta were incubated with DMEM supplemented with 10% FBS overnight at 37°C in the presence of 5% CO₂. The next day the aorta were cut into 2 mm segments and incubated with digestion buffer (175 U/mL of collagenase and 0.125 mg/mL of elastase in HBSS, filtered through 0.2 µm polyethersulfone membrane) for 1 hr at 37°C in the presence of 5% CO₂. Following digestion, the tissue was dissociated with a glass Pasteur pipette, DMEM supplemented with 10% FBS was added, and the cells were collected via centrifugation (200 × g for 8 min). The cells were re-suspended in DMEM supplemented with 20% FBS, transferred to a T-25 cm flask, and the cells were incubated overnight at 37°C in the presence of 5% CO₂. The next day the cells were carefully washed with DMEM supplemented with 10% FBS, 100 U/mL penicillin, and 100 µg/mL streptomycin and cultured normally with medium changes every three days. Cells were determined to be SMCs via immunofluorescence assays using the rabbit polyclonal smooth muscle α-actin antibody exactly as described by the manufacturer (1:1000, Abcam Inc., Cambridge, MA). For all experiments, mVSMCs were used between passages three and seven.

Isolation of Primary Cortical Neurons

Cortical neurons were prepared as previously detailed (34-36). Briefly, pups (postnatal day<2) were genotyped and euthanized by decapitation. The entire frontal cortex of RSK2^{+/+} and RSK2^{-/-} animals was microdissected from whole brain, followed by digestion in neurobasal medium containing 0.1% papain and 0.2% BSA at 37°C for 20 min. The medium was then replaced with complete neurobasal medium and the digested tissue was mechanically dissociated via trituration with a glass Pasteur pipette. The supernatant was transferred to a new sterile 1.5 mL tube and cells were collected via centrifugation (200 × g for 5 min). The cell pellet was then resuspended in conditioned complete neurobasal medium. Cells were counted and seeded at a density of 50,000 cells/well onto poly-L-lysine-coated 96 well plates (0.1 mg/mL low molecular mass poly-L-lysine, 0.625% boric acid, 0.955% sodium tetraborate) and cultured normally.

Lentivirus Production and Infection

Lentiviral infection of primary cortical neurons was performed essentially as previously described (35). In brief, a pre-determined mixture of 5-HT_{2A}-GFP-CT-FUGW and the viral packaging constructs VSVG and Delta 8.9 (ratio= 3.3 FUGW: 2.5 Delta 8.9 : 1 VSVG) were co-transfected into HEK293T cells using Fugene6. Forty-eight hours after transfection the medium containing virus was removed, pooled, and a virus pellet was obtained via centrifugation (26,000 × g for 5 hr). The virus pellet was re-suspended in PBS, concentrated approximately 40-fold using Amicon UltraCel 100K filters (Millipore), and then tested for infection and expression of 5-HT_{2A}-GFP in HEK293T cells. A pre-determined amount of concentrated lentivirus was then applied to primary cortical neurons cultured for 7-10 days *in vitro*.

Immunoprecipitation and Western Blotting

Immunoprecipitation of RSK2 and detection of Ser(P)-386 following growth factor treatment was performed as previously described (37). Briefly, RSK2^{+/+} and RSK2^{-/-} cells were treated with EGF (100ng/mL) or IGF-1 (10nM) for various times and solubilized with cold lysis buffer (50 mM Tris-HCl, 150mM NaCl, 1% tergitol, 0.5% sodium deoxycholate, 0.1% sodium dodecyl sulfate, EDTA-free protease inhibitors, 50mM NaF, 50mM β-glycerolphosphate, 5mM sodium pyrophosphate, and 0.1mM sodium orthovanadate, pH 8.0) for 20 min at 4°C. Supernatants were collected via centrifugation (15,000 × g, 30 min) and equal amounts of Protein A/G-cleared lysate were incubated with mouse monoclonal anti-RSK2 (2μg) for 2 hr at 4°C, followed by incubation with Protein A/G agarose for 2 hr at 4°C. Immunopurified complexes were extensively washed with lysis buffer, eluted with 2× SDS sample buffer (125 mM Tris-HCl, 4% sodium dodecyl sulfate, 20% glycerol, 200 mM dithiothreitol, 0.2% Bromphenol Blue, pH 6.8), and stored at -80°C until further use.

Proteins were immunoblotted using standard procedures (28). Specifically, proteins were resolved on 10% SDS-PAGE gels, electroblotted onto nitrocellulose membranes (BioRad Laboratories, Hercules, CA), and blocked with standard blocking buffer (Tris-buffered saline, 0.1% Tween-20, and 5% nonfat dehydrated milk) for 1 hr at RT. Membranes were then incubated with primary antibodies diluted in standard blocking buffer or phospho-specific blocking buffer (5% BSA and TBST). Specifically, RSK2 was detected using the goat polyclonal RSK2 antibody (1:1000, Santa Cruz Biotechnology, Inc.) and the rabbit polyclonal Ser(P)-386 antibody (1:1000, Cell Signaling Technology, Inc., Danvers, MA). The EGFR was detected using the rabbit polyclonal EGFR antibody (1:500, Santa Cruz Biotechnology, Inc.) and the mouse monoclonal Tyr(P)-1068 antibody (1:500, Cell Signaling Technology, Inc.). Phosphorylated IGF-1 R was detected using the rabbit polyclonal Tyr(P)-1158/1162/1163 antibody (1:1000, Upstate Millipore). Membranes were washed extensively with Tris-buffered saline + 0.1% Tween-20 (TBST) and subsequently incubated for 1 hr at RT with secondary horseradish peroxidase-conjugated antibodies raised against mouse, rabbit and goat IgG (1:1000, Vector Laboratories, Burlingame, CA). Membranes were washed extensively and proteins were detected using SuperSignal West Pico chemiluminescent substrate (Thermo Fisher Scientific, Inc., Rockford, IL). Immunoreactive bands were imaged and quantified using Kodak Imaging software (Eastman Kodak, New Haven, CT). Sum pixel intensity values were analyzed via the one-tailed, paired t test (significance defined as $p < 0.05$) (Graphpad Software, Inc., La Jolla, CA).

Fluorometric Imaging Plate Reader (FLIPR^{Tetra}) Analysis of Intracellular Ca²⁺ Release

Intracellular Ca²⁺ release was measured in MEFs and mVSMCs via FLIPR^{Tetra} assays using a Ca²⁺ assay kit (Molecular Devices, Sunnyvale, CA) as previously detailed (13, 14). Briefly, MEFs were plated at a density of 25,000 cells/well into black-wall, clear-bottom 96-

well tissue culture plates (Greiner Bio-One, Monroe, NC), whereas mVSMCs were plated at a density of 10,000 cells/well into black-wall, clear-bottom 384-well tissue culture plates (Greiner Bio-One). The cells were cultured in dialyzed medium (DMEM, 5% FBS dialyzed to <0.05 nM 5-HT, 1 mM sodium pyruvate, 100 units/mL penicillin, and 100 µg/mL streptomycin) and serum-free medium (DMEM, 0.1% BSA, 100 units/mL penicillin, and 100 µg/mL streptomycin) for 24-40 hr before the assay for normal and growth factor desensitization experiments, respectively. For both experiments, the cells were incubated with Ca^{2+} assay buffer (20 mM HEPES, 1× HBSS, 2.5 mM probenecid, and Ca^{2+} assay reagent, pH 7.4) for 60 min at 37°C prior to initiating the FLIPR program. However, for growth factor desensitization experiments the cells were incubated with growth factors diluted in Ca^{2+} assay buffer (for 30 min and 60 min time points). After dye loading, the FLIPRTetra was programmed to add agonist approximately 10 seconds after establishing baseline relative fluorescence unit (RFU) values (excitation 470-495, emission 515-575 nm). RFU values were collected every second for 5 min and the average baseline values were subtracted from maximum RFU values. Values were expressed relative to the maximal untreated response in each cell line and analyzed by nonlinear regression to generate fit parameters of potency (EC_{50}) and maximal signaling (E_{max}) (Graphpad software). The F test was used to determine the statistical significance (defined as $p < 0.05$) of the fit parameters in growth factor-treated vs. untreated cells.

Analysis of Intracellular Ca^{2+} Release in Primary Cortical Neurons

Cortical neurons were isolated, cultured and imaged as described previously (35, 36). In brief, 48 hr after lentivirus infection each well was imaged for total GFP fluorescence using the BD Pathway 855 high content imaging microscope equipped with environmental control. Ca^{2+} flux was then determined using the FLIPR Ca^{2+} assay kit (Molecular Probes) as detailed by the manufacturer. In brief, prior to live cell imaging, cells were washed 1× with phosphate buffered saline followed by 60 min incubation with Ca^{2+} assay buffer (20 mM HEPES, 1× HBSS, 2.5 mM probenecid, 0.57 mM ascorbic acid, and Ca^{2+} assay reagent, pH 7.4). Assays using growth factors were performed similarly except that growth factor was added during the dye loading step. Cells were maintained at 37°C during the entire period of observation and were imaged for 20 sec prior to drug addition to obtain baseline dye fluorescence. The liquid handling capability of the BD Pathway 855 was used to add 10× drug and then fluorescence images were obtained for 120 sec. To control for subtle differences in receptor expression, Ca^{2+} responses were normalized to GFP intensity/well using custom written macros for Excel (Microsoft, Redmond, WA) and Image J (U. S. National Institutes of Health, Bethesda, Maryland). Values were expressed as fold over baseline and a two-tailed, paired t test was used to determine the statistical significance (defined as $p < 0.05$) of responses in growth factor-treated vs. untreated cells.

Results

RSK2 is required for EGF-induced attenuation of 5-HT_{2A} receptor signaling

We recently determined that RSK2 interacts with the 5-HT_{2A} serotonin receptor and attenuates signaling *via* direct receptor phosphorylation (13, 14). Moreover, preliminary data suggested that EGFR activation attenuates 5-HT_{2A} receptor signaling, presumably by activating RSK2. Considering the potential for describing how 5-HT_{2A} receptors are regulated in cells, and perhaps *in vivo*, we applied pharmacological and genetic approaches to determine if various RTKs, including members of the EGFR family, require RSK2 for attenuating 5-HT_{2A} receptor signaling.

First, we activated the EGFR (also known as ErbB1) in RSK2^{+/+} and RSK2^{-/-} MEFs with two canonical agonists (i.e., EGF and TGF- α) and monitored 5-HT_{2A} receptor

responsiveness using 5-HT_{2A} agonists of varying intrinsic efficacies. Consistent with our initial EGFR findings (14), EGF significantly attenuated 5-HT_{2A} receptor signaling in RSK2^{+/+} MEFs. As shown in Figure 1A, 100ng/mL EGF maximally activated the EGFR and RSK2 in these cells. When receptor signaling was assayed, we found that EGF pretreatment resulted in significant rightward shifts in 5-HT concentration-response curves (CRCs) as early as 30min, with maximal effects reached within 60 min (Figure 1B, Table 1). According to classical concepts of receptor pharmacology, these rightward shifts in 5-HT CRCs denoted decreases in agonist potency, most likely resulting from attenuation of receptor signaling given the short time scale of the experiment (38). To best illustrate this decrease in 5-HT potency, we compared the 5-HT_{2A}-mediated Ca²⁺ responses elicited by a sub-maximal, EC₅₀ concentration of 5-HT (i.e., 10nM). As shown in Figure 1C, 10nM 5-HT elicited significantly lower Ca²⁺ responses after treating with EGF for 30 min (peak Ca²⁺ release= 99.7 ± 0.1% vs. 79.7 ± 3.1% in untreated and EGF-treated RSK2^{+/+} MEFs, respectively; N=3 to 9, p<0.05) and 60 min (peak Ca²⁺ release= 99.7 ± 0.08% vs. 63.7 ± 6.0% in untreated and EGF-treated RSK2^{+/+} MEFs, respectively; N=7 to 9, p<0.05). Although it was clear that EGF decreased 5-HT signaling, these results could be explained by non-specific effects associated with a single, supramaximal concentration of EGF. To address this concern we determined that EGF attenuated 5-HT_{2A} receptor signaling with an IC₅₀ of 1.3 ng/mL (Figure 1D), a value that is within the concentration range typically observed for EGFR-mediated signaling events (39).

In contrast to these results in RSK2^{+/+} MEFs, EGF cannot attenuate 5-HT_{2A} receptor signaling in RSK2^{-/-} MEFs. Specifically, EGF did not significantly alter 5-HT potency in RSK2^{-/-} MEFs as evidenced by superimposed 5-HT CRCs (Figure 1E and Table 1). Moreover, 5-HT_{2A}-mediated Ca²⁺ responses elicited by an EC₅₀ concentration of 5-HT were not significantly decreased after treating with EGF for 30 min (peak Ca²⁺ release= 99.7 ± 0.2% vs. 93.9 ± 4.2% in untreated and EGF-treated RSK2^{-/-} MEFs, respectively; N=3 to 6, p>0.05) and 60 min (peak Ca²⁺ release= 99.7 ± 0.2% vs. 96.8 ± 6.3% in untreated and EGF-treated RSK2^{-/-} MEFs, respectively; N=6, p>0.05) (Figure 1F).

It was conceivable that differences in EGFR signal transduction between RSK2^{+/+} and RSK2^{-/-} MEFs could account for the differential effects of EGF. To evaluate this possibility, we compared the mRNA expression profiles of genes constituting the EGFR signal transduction pathway in RSK2^{+/+} and RSK2^{-/-} MEFs. As shown in Figure 2, analysis of microarray studies revealed no substantial differences in gene expression profiles between RSK2^{+/+} and RSK2^{-/-} MEFs that could account for lack of attenuation in RSK2^{-/-} MEFs. Only two differences were apparent in RSK2^{-/-} cells: (1) a decrease in RSK2 mRNA (as predicted in knock-out cells) and (2) an increase in Sos2 mRNA. Additionally, at the protein level we found that EGFR activation was similar between RSK2^{+/+} and RSK2^{-/-} MEFs after 5 min of EGF treatment (472 ± 131% vs. 491 ± 96% for EGFR phosphorylation in RSK2^{+/+} and RSK2^{-/-} MEFs, respectively; N=3, p>0.05) (Figure 1A). Taken together, these findings indicate (1) that RSK2 is a critical mediator of inhibitory cross-talk between EGF and 5-HT_{2A} receptors in MEFs and (2) that the effects are not due to compensatory changes in expression of EGFR signaling partners.

We further confirmed that RSK2 was required for inhibitory cross-talk by activating the EGFR in RSK2^{+/+} and RSK2^{-/-} MEFs with another selective and potent EGFR agonist, TGF- α . Similar to our results with EGF, TGF- α attenuated 5-HT_{2A} receptor signaling in RSK2^{+/+} MEFs. Specifically, 1hr treatment with TGF- α decreased 5-HT potency, as illustrated by significant rightward shifts in 5-HT CRCs (Figure 3A and Table 1) and significant decreases in 5-HT_{2A}-mediated Ca²⁺ release elicited by an EC₅₀ concentration of 5-HT (peak Ca²⁺ release= 99.7 ± 0.2% vs. 76.9 ± 2.2% in untreated and TGF- α -treated

RSK2^{+/+} MEFs, respectively; N=5, $p<0.05$) (Figure 3B). Moreover, TGF- α attenuated 5-HT_{2A} receptor signaling in RSK2^{+/+} MEFs with an IC₅₀ of 4.1 ng/mL (Figure 3C), again consistent with EGFR-mediated signaling events. In agreement with our previous experiments using EGF, TGF- α treatment did not attenuate 5-HT_{2A} receptor signaling in RSK2^{-/-} MEFs (Figure 3D and Table 1). Moreover, we did not detect large decreases in 5-HT_{2A}-mediated Ca²⁺ release elicited by an EC₅₀ concentration of 5-HT (peak Ca²⁺ release= 99.8 \pm 0.2% vs. 91.7 \pm 1.2% in untreated and TGF- α -treated RSK2^{-/-} MEFs, respectively; $p<0.05$) (Figure 3E).

In addition to full agonists, partial agonists display characteristic and predictable signaling behaviors under conditions of decreased receptor responsiveness. Notably, in this context, full agonists commonly exhibit effects on potency but lesser effects on maximal signaling, while partial agonists commonly display decreases in maximal signaling (i.e., large downward shifts in CRCs) (38). Thus, we predicted that EGF treatment would decrease the maximal signaling of the weak partial agonist lisuride with minimal effects on potency; whereas the potency of the strong partial agonist 5-methoxyDMT would be significantly decreased. Consistent with these predictions, EGF treatment significantly decreased the maximal signaling of lisuride in RSK2^{+/+} MEFs (Figure 4A, Table 1). Furthermore, 5-methoxyDMT displayed behaviors intermediate between the full agonist 5-HT and the weak partial agonist lisuride, exhibiting a minor decrease in maximal signaling and a significant decrease in potency (Figure 4C, Table 1). In agreement with a requirement for RSK2, we did not observe significant shifts in the CRCs of either partial agonist in RSK2^{-/-} MEFs (Figures 4B and 4D, Table 1). Taken together, these pharmacological and genetic approaches strongly support the hypothesis that EGFRs act via RSK2 to attenuate 5-HT_{2A} receptor signaling in MEFs.

RSK2 is required for PDGFR-mediated attenuation of endogenous 5-HT_{2A} receptor signaling in primary mVSMCs

The growth factor PDGF is a potent mitogen, chemoattractant, and survival factor that activates RSKs downstream of PDGFR activation in VSMCs (40, 41). VSMCs also endogenously express 5-HT_{2A} receptors which produce measurable Ca²⁺ responses in FLIPR assays (Figure 5A) (42). Therefore, VSMCs isolated from RSK2^{+/+} and RSK2^{-/-} mice represented an intact model system whereby we could test: (1) whether inhibitory cross-talk occurs between additional RTKs and endogenously expressed 5-HT_{2A} receptors, and (2) to what extent this requires RSK2.

In these studies we activated PDGFRs with PDGF-AB and PDGF-BB, the principal PDGF ligands in serum (43). As evidenced by significant downward shifts in 5-HT CRCs following 60 min treatments with PDGF-AB (Figure 5B) and PDGF-BB (Figure 5D), activation of the PDGFR resulted in attenuation of 5-HT_{2A} receptor signaling (Table 2). To best illustrate this we showed that PDGF-BB treatment significantly decreased 5-HT_{2A}-mediated Ca²⁺ responses in RSK2^{+/+} MEFs elicited by a saturating concentration of 5-HT (i.e., 10 μ M) (peak Ca²⁺ release= 90.2 \pm 0.7% vs. 67.0 \pm 8.5% in untreated and PDGF-BB-treated RSK2^{+/+} MEFs, respectively; N=4, $p<0.05$) (Figure 5E). As expected, PDGF treatments did not significantly reduce the maximal signaling of 5-HT in RSK2^{-/-} mVSMCs (Figures 5C and 5F). Moreover, PDGF-BB treatment failed to significantly decrease 5-HT_{2A}-mediated Ca²⁺ responses in RSK2^{-/-} MEFs elicited by a saturating concentration of 5-HT (peak Ca²⁺ release= 90.3 \pm 3.8% vs. 77.9 \pm 7.6% in untreated and PDGF-BB-treated RSK2^{-/-} MEFs, respectively; N=4, $p>0.05$) (Figure 5G). Together with our results using two different EGFR agonists, these results strongly suggest that RSK2 is required for inhibitory cross-talk between multiple growth factor signaling pathways and the 5-HT_{2A} receptor.

IGF-1 weakly activates RSK2 in MEFs and does not attenuate 5-HT_{2A} receptor signaling

We have demonstrated using various RTK agonists (i.e. EGF, TGF- α , and PDGF), cell lines (MEFs and VSMCs), and GPCR ligands (i.e. 5-HT, 5-methoxy-DMT, and lisuride) that RTKs require RSK2 to attenuate 5-HT_{2A} receptor signaling. However, it was unknown whether insulin or IGF-1, which have been shown to attenuate the signaling of GPCRs including the closely related 5-HT_{2C} receptor (22), also attenuate 5-HT_{2A} receptor signaling.

In initial experiments testing insulin, we determined that insulin showed no effect on 5-HT_{2A} receptor signaling, despite modest activation of RSK2 (data not shown). However, upon closer examination we discovered that insulin receptors are not expressed at detectable levels in RSK2^{+/+} and RSK2^{-/-} MEFs (Figure S1), suggesting that RSK2 activation is mediated via the IGF-1 R which has low affinity for insulin (44). Since both RSK2^{+/+} and RSK2^{-/-} MEFs express equal amounts of IGF-1 R (Figure S1), we next determined if IGF-1 could attenuate 5-HT_{2A} receptor signaling. As shown in Figure 6A and 6B, IGF-1 treatment did not result in large shifts in 5-HT CRCs in RSK2^{+/+} or RSK2^{-/-} MEFs (Table 1), identical to our results with insulin. Importantly, these results could not be explained by a general deficiency in RTK signaling since a 1hr treatment with EGF attenuated 5-HT_{2A} signaling in parallel control experiments (peak Ca²⁺ release= 99.9 \pm 0.1% vs. 80.7 \pm 2.3% in untreated and EGF-treated RSK2^{+/+} MEFs, respectively; N=3) (Figure 6C). Additional experimentation showed that the IGF-1R is activated by IGF-1 (Figure S2). Together, these findings suggested that the mechanism(s) underlying inhibitory cross-talk between an RTK and a GPCR (e.g., RSK2 activation) engenders some level of specificity. One such possibility was that IGF-1 treatment only modestly activated RSK2 (24). Indeed, as shown in Figure 6D and quantified in Figure 6E, maximal activation of RSK2 by IGF-1 was significantly less when compared to EGF ($0.546 \times 10^6 \pm 0.107 \times 10^6$ vs. $1.25 \times 10^6 \pm 0.25 \times 10^6$ for IGF-1 and EGF, respectively; N=3, $p < 0.05$). Thus, robust activation of RSK2 by RTKs seems to be required for inhibitory cross-talk with the 5-HT_{2A} receptor. Regardless of the mechanism, it is likely that some degree of specificity exists given the potential physiological importance of RTK-GPCR crosstalk.

RSK2 is required for EGF-mediated attenuation of endogenous P2Y purinergic receptor signaling

RSK2 attenuates the signaling of additional GPCRs endogenously expressed in MEFs, including P2Y purinergic receptors (13). Therefore, we hypothesized that EGFR activation, in addition to regulating 5-HT_{2A} receptors, could also attenuate P2Y receptor signaling in a RSK2-dependent manner. By testing this hypothesis we could begin to address whether this novel regulatory mechanism is conserved across multiple GPCRs that show sensitivity to RSK2 regulation.

As shown in Figure 7A, EGF treatment significantly reduced ATP signaling in RSK2^{+/+} MEFs (Table 1). Specifically, we observed significant decreases in ATP maximal signaling (E_{\max} =99.7 \pm 2.0% vs. 82.5 \pm 3.2% in untreated and EGF-treated RSK2^{+/+} MEFs, respectively; N=5, $p < 0.05$) and potency (7.0 μ M vs. 13 μ M for untreated and hEGF-treated, respectively; N=5, $p < 0.05$) following 60 min EGF treatment. Similar to our observations in mVSMCs (Figure 5), decreased maximal signaling of the full agonist ATP was consistent with desensitization of endogenously expressed P2Y receptors. However, treating RSK2^{-/-} MEFs with EGF failed to significantly decrease ATP maximal signaling or potency (Figure 7B, Table 1). These data are important because they provide the first evidence for a common regulatory mechanism whereby RTKs act via RSK2 to regulate the signaling of multiple GPCRs.

Growth factors essential for normal brain function attenuate 5-HT_{2A} receptor signaling in cortical neurons

We have presented multiple lines of evidence to show that 5-HT_{2A} signaling is indeed attenuated following activation of several endogenous RTKs in multiple cell types. In addition to expression in peripheral tissues, RTKs are widely expressed throughout the brain (e.g., in the cortex) and are activated by endogenous ligands such as EGF and NRG-1 (45). Since 5-HT_{2A} receptors are also highly expressed in the cortex (46), it was tempting to speculate that cross-talk between RTKs and 5-HT_{2A} receptors could explain how 5-HT_{2A} receptors are regulated in cortical neurons.

To test this possibility, we developed a live cell imaging technique to measure 5-HT_{2A} receptor signaling in cortical neurons in the presence and absence of growth factors. As shown in Figure 8A and 8B, uninfected neurons were unresponsive to the 5-HT_{2A/2C} selective agonist DOI, despite robust Ca²⁺ responses following depolarization with 80 mM KCl (Figure 8C). However, DOI elicited measurable Ca²⁺ responses in neurons only after infection with GFP-tagged 5-HT_{2A} receptors (Figure 8D and 8E), thus ensuring specificity of the DOI response. We then quantified these DOI-induced responses in untreated (Figure 8E) and growth factor-treated (Figure 8H) neurons via manual segmenting (Figure 8F and 8I). As shown in Figure 8J and quantified in Table 3, treatment with either EGF or NRG-1 significantly reduced the Ca²⁺ response elicited by DOI. In these studies we present the first evidence that inhibitory cross-talk occurs between RTKs and GPCRs in neurons. Most importantly, our data show that RTKs attenuate 5-HT_{2A} signaling in neurons—a finding with enormous potential for explaining how 5-HT_{2A} receptors are regulated in the brain.

Discussion

The three major findings in this paper are: 1) multiple endogenous RTK receptors and their ligands attenuate 5-HT_{2A} receptor responsiveness in several physiologically relevant cell types; 2) RSK2 is required for RTK-mediated attenuation of 5-HT_{2A} receptor signaling, and 3) RTK activation similarly attenuates P2Y purinergic signaling in a RSK2-dependent manner. By directly testing multiple endogenous growth factors/RTK pathways and multiple Gq-coupled GPCRs, we have now established a cellular mechanism whereby RTK signaling cascades attenuate GPCR signaling through RSK2. Importantly, these findings support a novel paradigm of inhibitory cross-talk between RTKs and GPCRs and extend it to include a larger mechanism whereby RTKs act via RSK2 to regulate the signaling of multiple GPCRs.

RSK2 is required for inhibitory cross-talk between RTKs and the 5-HT_{2A} receptor in a variety of cell types

Consistent with evidence for inhibitory cross-talk between RTKs and select GPCRs (i.e., the β_1 -, β_2 -, α_{1B} -, and α_{1D} -adrenergic receptors, and 5-HT_{2C} receptor) (16-22), our data demonstrate that activation of the EGFR attenuates 5-HT_{2A} receptor signaling in MEFs, VSMCs and cortical pyramidal neurons. Moreover, we discovered that this novel regulatory pathway requires RSK2. We verified that the EGFR requires RSK2 to attenuate 5-HT_{2A} receptor signaling by observing the signaling of both full and partial 5-HT_{2A} agonists in RSK2^{+/+} and RSK2^{-/-} cells. Since changes to GPCR responsiveness affect each agonist class differently, this approach allowed us to unambiguously identify RTK-mediated effects on receptor signaling. Explicitly, full agonists have a large receptor reserve and are resistant to changes in the population of functional receptors (i.e., resulting from receptor desensitization or down-regulation). As a result, full agonists signal maximally but with lower potency under conditions of receptor desensitization in cells over-expressing a GPCR (i.e., CRCs are right-shifted) (38). However, both the maximal signaling and potency of full

agonists is decreased under conditions of receptor desensitization in cells with endogenous GPCR expression (i.e., CRCs are predominantly shifted downward with minor rightward shifts). Partial agonists, on the other hand, have low receptor reserve and are more sensitive to changes in the population of functional GPCRs. As a result, partial agonists typically signal with lower efficacy and potency under conditions of receptor desensitization irrespective of receptor expression (38). In line with these predictions, we observed that EGF significantly decreased full agonist (i.e., 5-HT) potency and partial agonist (i.e., lisuride) efficacy in high-expressing RSK2^{+/+} MEFs. Additionally, growth factor treatment significantly decreased the maximal signaling of full agonists (i.e., 5-HT and ATP) when their cognate receptors were expressed at endogenous levels in RSK2^{+/+} mVSMCs and MEFs. Taken together, our results in RSK2^{+/+} cells are consistent with RTK-mediated attenuation of 5-HT_{2A} receptor signaling. Importantly, none of these predicted effects were observed in RSK2^{-/-} MEFs, thus supporting the hypothesis that RTKs act via RSK2 to attenuate 5-HT_{2A} receptor signaling.

Alternatively, these results could be explained by differences in gene expression profiles between RSK2^{+/+} and RSK2^{-/-} cells. However, microarray data show that the expression of genes required for EGFR signal transduction are not significantly different between RSK2^{+/+} and RSK2^{-/-} MEFs. Thus, the simplest explanation for our results remains that RSK2 is a critical mediator of cross-talk between the EGFR and 5-HT_{2A} receptor.

In addition to our results in RSK2^{+/+} MEFs, we observed that 5-HT_{2A} signaling was significantly decreased following activation of endogenous RTKs in mVSMC and cortical neuron primary cell lines. Considering the physiological importance of inhibitory cross-talk, it is attractive to speculate that growth factor signaling may be relevant for regulating the 5-HT_{2A} receptor in the CNS. In support of this, members of the EGFR family are widely expressed throughout the brain and regulate a variety of functions including proliferation, differentiation, maturation, and survival of a variety of neurons (45). Interestingly, the ErbB4 neuregulin receptor, which is a member of the EGFR family, is expressed throughout the mature brain and is known to reside in some of the same cortical layers (47) as the 5-HT_{2A} receptor (48). Moreover, ErbB4 interacts with PSD-95, a post-synaptic density protein that associates with and regulates 5-HT_{2A} receptor signaling and trafficking *in vitro* and *in vivo* (26, 29, 35, 49). Thus, considering the pervasiveness of growth factor signaling in the brain, as well as its overlapping expression with 5-HT_{2A} receptors, RTK signaling could modulate 5-HT_{2A} receptor signaling *in vivo*. Intriguingly, aberrant signaling of both RTKs and 5-HT_{2A} receptors has been associated with neuropsychiatric disorders such as depression and schizophrenia (47, 50-52). Together, these findings suggest that a more complete understanding of the mechanism(s) underlying inhibitory cross-talk between RTKs and 5-HT_{2A} receptors is of considerable therapeutic importance.

IGF-1 fails to robustly activate RSK2 and does not attenuate 5-HT_{2A} receptor signaling

In stark contrast to our results using EGF, PDGF, and ErbB4 receptor agonists, we discovered that IGF-1 did not attenuate 5-HT_{2A} receptor signaling in either RSK2^{+/+} or RSK2^{-/-} MEFs. In order to interpret these negative results, we showed that EGF treatment retained the ability to attenuate 5-HT_{2A}-mediated Ca²⁺ release in parallel control experiments. These data suggest that, unlike EGF receptor activation, IGF-1 signaling does not desensitize 5-HT_{2A} receptors. The reasons for this are unknown, although our data showing that IGF-1 weakly activates RSK2 when compared to EGF suggest that a threshold level of RSK2 activation must be reached in order to elicit 5-HT_{2A} receptor desensitization.

Other mechanisms have been proposed to explain IGF-1-induced GPCR desensitization including phosphorylation of tyrosine residues in the second intracellular loop of the β_2 -adrenergic receptor and Akt-mediated phosphorylation of the β_1 -adrenergic receptor (16, 19). However, 5-HT_{2A} receptors are not known to be phosphorylated on tyrosine residues and are not substrates for Akt, perhaps explaining why IGF-1 has no effect on 5-HT_{2A} receptor signaling.

RSK2 is required for growth factor-mediated regulation of multiple GPCRs-evidence from P2Y purinergic receptors

In addition to the 5-HT_{2A} receptor, endogenous P2Y purinergic receptor signaling is regulated by RSK2 (13). Here we provide the first evidence showing that, like 5-HT_{2A} receptors, EGFR activation attenuates P2Y-purinergic receptor signaling in a RSK2-dependent manner. Thus, it appears that RSK2 is a critical mediator of inhibitory crosstalk between multiple RTKs and GPCRs. Interestingly, the β_1 -adrenergic and PAR-1 thrombinergic receptors are also regulated by RSK2 (13), and it remains to be determined if these receptors are regulated by RTKs in a RSK2-dependent manner. This is an especially intriguing question for the β_1 AR since it is already known that activation of the IGF-1R regulates β_1 -adrenergic receptor signaling through activation of PI3 kinase and Akt (16).

A question of important physiological relevance is whether specific RTK signaling pathways influence the signaling of all or only select groups of GPCRs. Our results, along with those of others, indicate that signaling from some Gq-coupled receptors (i.e., 5HT_{2A}, P2Y, α_{1b} -adrenergic, and α_{1d} -adrenergic) are similarly attenuated by one RTK, the EGFR (20, 21). Other RTKs, such as insulin and IGF-1 receptors, are well-known to decrease the signaling of some Gs-coupled GPCRs such as β_1 - and β_2 -adrenergic receptors (16, 53). However, insulin and IGF-1 receptors attenuate signaling from only some (i.e., 5-HT_{2C}), but not all (i.e., M1 muscarinic or 5-HT_{2A}) Gq-coupled GPCRs ((22), Figure 6). Therefore, a robustness of this signaling crosstalk is evident and RTK inhibitory crosstalk to GPCRs will likely emerge as a receptor-specific phenomenon. Ultimately, further studies testing many RTKs and GPCRs will help elucidate if this crosstalk is a conserved phenomenon.

In summary, multiple lines of evidence suggest that RSK2 is a critical mediator of inhibitory cross-talk between RTKs and the 5-HT_{2A} receptor. Specifically, this study presents the first evidence that 5-HT_{2A} receptor signaling is attenuated by the growth-factor-mediated activation of RTKs endogenously expressed in multiple cell types including physiologically relevant mVSMCs and cortical neurons. Moreover, genetic deletion of RSK2 was sufficient to block these effects, thus demonstrating that RSK2 is required for the inhibitory cross-talk between RTKs and 5-HT_{2A} receptors in all relevant cell types examined. Intriguingly, we discovered that the P2Y purinergic receptor, whose signaling is also regulated by RSK2, is similarly attenuated following EGFR activation in RSK2^{+/+} MEFs. Taken together, these findings provide the initial framework for a conserved regulatory mechanism whereby multiple RTKs act via the ERK/MAPK effector RSK2 to attenuate GPCR signaling. Most importantly, inhibitory cross-talk between RTKs and 5-HT_{2A} receptors could provide insight into how these receptors are regulated *in vivo*.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Kimberly Molnar for her expert assistance with mVSMC isolation and Dr. Betsy Pehek for HPLC-electrochemical detection analysis of dialyzed serum.

Abbreviations

5-HT	serotonin
5-HT_{2A}	serotonin 2A receptor
GPCR	G protein-coupled receptor
RSK2	p90 ribosomal S6 kinase 2
ERK/MAPK	extracellular signal regulated kinase/mitogen activated protein kinase
CRC	concentration-response curve
RTK	receptor tyrosine kinase
EGF	epidermal growth factor
EGFR	epidermal growth factor receptor
IGF-1	insulin-like growth factor 1
IGF-1 R	insulin-like growth factor 1 receptor
TGF-α	transforming growth factor-alpha
PDGF	platelet-derived growth factor
PDGFR	platelet-derived growth factor receptor
NRG-1	neuregulin-1
GFP	green fluorescent protein
5-methoxyDMT	5-methoxy-N,N-dimethyltryptamine
DOI	(\pm)-2,5-dimethoxy-4-iodoamphetamine hydrochloride
FBS	fetal bovine serum
DMEM	Dulbecco's modified Eagle's medium
HBSS	Hank's buffered salt solution
MEF	mouse embryonic fibroblast
mVSMC	mouse vascular smooth muscle cell
RFU	relative fluorescent unit

References

1. King N, Hittinger CT, Carroll SB. Evolution of key cell signaling and adhesion protein families predates animal origins. *Science*. 2003; 301:361–363. [PubMed: 12869759]
2. Schioth HB, Nordstrom KJ, Fredriksson R. Mining the gene repertoire and ESTs for G protein-coupled receptors with evolutionary perspective. *Acta Physiol (Oxf)*. 2007; 190:21–31. [PubMed: 17428229]
3. Venter JC, Adams MD, Myers EW, Li PW, Mural RJ, Sutton GG, Smith HO, Yandell M, Evans CA, Holt RA, Gocayne JD, Amanatides P, Ballegu RM, Huson DH, Wortman JR, Zhang Q, Kodira CD, Zheng XH, Chen L, Skupski M, Subramanian G, Thomas PD, Zhang J, Miklos G. L. Gabor, Nelson C, Broder S, Clark AG, Nadeau J, McKusick VA, Zinder N, Levine AJ, Roberts RJ, Simon M, Slayman C, Hunkapiller M, Bolanos R, Delcher A, Dew I, Fasulo D, Flanigan M, Florea L,


Halpern A, Hannenhalli S, Kravitz S, Levy S, Mobarry C, Reinert K, Remington K, Abu-Threideh J, Beasley E, Biddick K, Bonazzi V, Brandon R, Cargill M, Chandramouliswaran I, Charlab R, Chaturvedi K, Deng Z, Di Francesco V, Dunn P, Eilbeck K, Evangelista C, Gabrielian AE, Gan W, Ge W, Gong F, Gu Z, Guan P, Heiman TJ, Higgins ME, Ji RR, Ke Z, Ketchum KA, Lai Z, Lei Y, Li Z, Li J, Liang Y, Lin X, Lu F, Merkulov GV, Milshina N, Moore HM, Naik AK, Narayan VA, Neelam B, Nusskern D, Rusch DB, Salzberg S, Shao W, Shue B, Sun J, Wang Z, Wang A, Wang X, Wang J, Wei M, Wides R, Xiao C, Yan C, Yao A, Ye J, Zhan M, Zhang W, Zhang H, Zhao Q, Zheng L, Zhong F, Zhong W, Zhu S, Zhao S, Gilbert D, Baumhueter S, Spier G, Carter C, Cravchik A, Woodage T, Ali F, An H, Awe A, Baldwin D, Baden H, Barnstead M, Barrow I, Beeson K, Busam D, Carver A, Center A, Cheng ML, Curry L, Danaher S, Davenport L, Desilets R, Dietz S, Dodson K, Doup L, Ferriera S, Garg N, Gluecksmann A, Hart B, Haynes J, Haynes C, Heiner C, Hladun S, Hostin D, Houck J, Howland T, Ibegwam C, Johnson J, Kalush F, Kline L, Koduru S, Love A, Mann F, May D, McCawley S, McIntosh T, McMullen I, Moy M, Moy L, Murphy B, Nelson K, Pfannkoch C, Pratts E, Puri V, Qureshi H, Reardon M, Rodriguez R, Rogers YH, Romblad D, Ruhfel B, Scott R, Sitter C, Smallwood M, Stewart E, Strong R, Suh E, Thomas R, Tint NN, Tse S, Vech C, Wang G, Wetter J, Williams S, Williams M, Windsor S, Winn-Deen E, Wolfe K, Zaveri J, Zaveri K, Abril JF, Guigo R, Campbell MJ, Sjolander KV, Karlak B, Kejariwal A, Mi H, Lazareva B, Hatton T, Narechania A, Diemer K, Muruganujan A, Guo N, Sato S, Bafna V, Istrail S, Lippert R, Schwartz R, Walenz B, Yooseph S, Allen D, Basu A, Baxendale J, Blick L, Caminha M, Carnes-Stine J, Caulk P, Chiang YH, Coyne M, Dahlke C, Mays A, Dombroski M, Donnelly M, Ely D, Esparham S, Fosler C, Gire H, Glanowski S, Glasser K, Glodek A, Gorokhov M, Graham K, Gropman B, Harris M, Heil J, Henderson S, Hoover J, Jennings D, Jordan C, Jordan J, Kasha J, Kagan L, Kraft C, Levitsky A, Lewis M, Liu X, Lopez J, Ma D, Majoros W, McDaniel J, Murphy S, Newman M, Nguyen T, Nguyen N, Nodell M, Pan S, Peck J, Peterson M, Rowe W, Sanders R, Scott J, Simpson M, Smith T, Sprague A, Stockwell T, Turner R, Venter E, Wang M, Wen M, Wu D, Wu M, Xia A, Zandieh A, Zhu X. The sequence of the human genome. *Science*. 2001; 291:1304–1351. [PubMed: 11181995]

4. Kristiansen K. Molecular mechanisms of ligand binding, signaling, and regulation within the superfamily of G-protein-coupled receptors: molecular modeling and mutagenesis approaches to receptor structure and function. *Pharmacol Ther*. 2004; 103:21–80. [PubMed: 15251227]
5. Kroeze WK, Sheffler DJ, Roth BL. G-protein-coupled receptors at a glance. *J Cell Sci*. 2003; 116:4867–4869. [PubMed: 14625380]
6. Thompson MD, Burnham WM, Cole DE. The G protein-coupled receptors: pharmacogenetics and disease. *Crit Rev Clin Lab Sci*. 2005; 42:311–392. [PubMed: 16281738]
7. Lagerstrom MC, Schioth HB. Structural diversity of G protein-coupled receptors and significance for drug discovery. *Nat Rev Drug Discov*. 2008; 7:339–357. [PubMed: 18382464]
8. Kroeze WK, Kristiansen K, Roth BL. Molecular biology of serotonin receptors structure and function at the molecular level. *Curr Top Med Chem*. 2002; 2:507–528. [PubMed: 12052191]
9. Berger M, Gray JA, Roth BL. The Expanded Biology of Serotonin. *Annu Rev Med*. 2009; 60:355–366. [PubMed: 19630576]
10. Nichols DE. Hallucinogens. *Pharmacol Ther*. 2004; 101:131–181. [PubMed: 14761703]
11. Roth BL, Baner K, Westkaemper R, Siebert D, Rice KC, Steinberg S, Ernsberger P, Rothman RB. Salvinorin A: a potent naturally occurring nonnitrogenous kappa opioid selective agonist. *Proc Natl Acad Sci U S A*. 2002; 99:11934–11939. [PubMed: 12192085]
12. Willins DL, Berry SA, Alsayegh L, Backstrom JR, Sanders-Bush E, Friedman L, Roth BL. Clozapine and other 5-hydroxytryptamine-2A receptor antagonists alter the subcellular distribution of 5-hydroxytryptamine-2A receptors in vitro and in vivo. *Neuroscience*. 1999; 91:599–606. [PubMed: 10366017]
13. Sheffler DJ, Kroeze WK, Garcia BG, Deutch AY, Hufeisen SJ, Leahy P, Bruning JC, Roth BL. p90 ribosomal S6 kinase 2 exerts a tonic brake on G protein-coupled receptor signaling. *Proc Natl Acad Sci U S A*. 2006; 103:4717–4722. [PubMed: 16537434]
14. Strachan RT, Sheffler DJ, Willard B, Kinter M, Kiselar JG, Roth BL. Ribosomal S6 kinase 2 directly phosphorylates the 5-hydroxytryptamine 2A (5-HT_{2A}) serotonin receptor, thereby modulating 5-HT_{2A} signaling. *J Biol Chem*. 2009; 284:5557–5573. [PubMed: 19103592]

15. Frodin M, Gammeltoft S. Role and regulation of 90 kDa ribosomal S6 kinase (RSK) in signal transduction. *Mol Cell Endocrinol.* 1999; 151:65–77. [PubMed: 10411321]
16. Gavi S, Yin D, Shumay E, Wang HY, Malbon CC. Insulin-like growth factor-I provokes functional antagonism and internalization of beta1-adrenergic receptors. *Endocrinology.* 2007; 148:2653–2662. [PubMed: 17363461]
17. Hadcock JR, Port JD, Gelman MS, Malbon CC. Cross-talk between tyrosine kinase and G-protein-linked receptors. Phosphorylation of beta 2-adrenergic receptors in response to insulin. *J Biol Chem.* 1992; 267:26017–26022. [PubMed: 1281480]
18. Karoor V, Baltensperger K, Paul H, Czech MP, Malbon CC. Phosphorylation of tyrosyl residues 350/354 of the beta-adrenergic receptor is obligatory for counterregulatory effects of insulin. *J Biol Chem.* 1995; 270:25305–25308. [PubMed: 7592686]
19. Karoor V, Malbon CC. Insulin-like growth factor receptor-1 stimulates phosphorylation of the beta2-adrenergic receptor in vivo on sites distinct from those phosphorylated in response to insulin. *J Biol Chem.* 1996; 271:29347–29352. [PubMed: 8910597]
20. Garcia-Sainz JA, Romero-Avila MT, Molina-Munoz T, Ldel C. Medina. Insulin induces alpha1B-adrenergic receptor phosphorylation and desensitization. *Life Sci.* 2004; 75:1937–1947. [PubMed: 15306161]
21. Rodriguez-Perez CE, Calvo-Ochoa E, Kalashnikova EV, Reyes-Cruz G, Romero-Avila MT, Garcia-Sainz JA. Receptor tyrosine kinases regulate alpha1D-adrenoceptor signaling properties: phosphorylation and desensitization. *Int J Biochem Cell Biol.* 2009; 41:1276–1283. [PubMed: 19038360]
22. Hurley JH, Zhang S, Bye LS, Marshall MS, DePaoli-Roach AA, Guan K, Fox AP, Yu L. Insulin signaling inhibits the 5-HT2C receptor in choroid plexus via MAP kinase. *BMC Neurosci.* 2003; 4:10. [PubMed: 12795815]
23. Hanauer A, Young ID. Coffin-Lowry syndrome: clinical and molecular features. *J Med Genet.* 2002; 39:705–713. [PubMed: 12362025]
24. Bruning JC, Gillette JA, Zhao Y, Bjorbaeck C, Kotzka J, Knebel B, Avci H, Hanstein B, Lingohr P, Moller DE, Krone W, Kahn CR, Muller-Wieland D. Ribosomal subunit kinase-2 is required for growth factor-stimulated transcription of the c-Fos gene. *Proc Natl Acad Sci U S A.* 2000; 97:2462–2467. [PubMed: 10716983]
25. Guan XM, Kobilka TS, Kobilka BK. Enhancement of membrane insertion and function in a type IIIb membrane protein following introduction of a cleavable signal peptide. *J Biol Chem.* 1992; 267:21995–21998. [PubMed: 1331042]
26. Xia Z, Gray JA, Compton-Toth BA, Roth BL. A direct interaction of PSD-95 with 5-HT2A serotonin receptors regulates receptor trafficking and signal transduction. *J Biol Chem.* 2003; 278:21901–21908. [PubMed: 12682061]
27. Morgenstern JP, Land H. A series of mammalian expression vectors and characterisation of their expression of a reporter gene in stably and transiently transfected cells. *Nucleic Acids Res.* 1990; 18:1068. [PubMed: 2156225]
28. Gray JA, Sheffler DJ, Bhatnagar A, Woods JA, Hufeisen SJ, Benovic JL, Roth BL. Cell-type specific effects of endocytosis inhibitors on 5-hydroxytryptamine(2A) receptor desensitization and resensitization reveal an arrestin-, GRK2-, and GRK5-independent mode of regulation in human embryonic kidney 293 cells. *Mol Pharmacol.* 2001; 60:1020–1030. [PubMed: 11641430]
29. Xia Z, Hufeisen SJ, Gray JA, Roth BL. The PDZ-binding domain is essential for the dendritic targeting of 5-HT2A serotonin receptors in cortical pyramidal neurons in vitro. *Neuroscience.* 2003; 122:907–920. [PubMed: 14643760]
30. Lois C, Hong EJ, Pease S, Brown EJ, Baltimore D. Germline transmission and tissue-specific expression of transgenes delivered by lentiviral vectors. *Science.* 2002; 295:868–872. [PubMed: 11786607]
31. Dahlquist KD, Salomonis N, Vranizan K, Lawlor SC, Conklin BR. GenMAPP, a new tool for viewing and analyzing microarray data on biological pathways. *Nat Genet.* 2002; 31:19–20. [PubMed: 11984561]


32. Doniger SW, Salomonis N, Dahlquist KD, Vranizan K, Lawlor SC, Conklin BR. MAPPFinder: using Gene Ontology and GenMAPP to create a global gene-expression profile from microarray data. *Genome Biol.* 2003; 4:R7. [PubMed: 12540299]
33. Moon SK, Thompson LJ, Madamanchi N, Ballinger S, Papaconstantinou J, Horaist C, Runge MS, Patterson C. Aging, oxidative responses, and proliferative capacity in cultured mouse aortic smooth muscle cells. *Am J Physiol Heart Circ Physiol.* 2001; 280:H2779–2788. [PubMed: 11356636]
34. Ahlemeyer B, Baumgart-Vogt E. Optimized protocols for the simultaneous preparation of primary neuronal cultures of the neocortex, hippocampus and cerebellum from individual newborn (P0.5) C57Bl/6J mice. *J Neurosci Methods.* 2005; 149:110–120. [PubMed: 16084598]
35. Abbas AI, Yadav PN, Yao WD, Arbuckle MI, Grant SG, Caron MG, Roth BL. PSD-95 is essential for hallucinogen and atypical antipsychotic drug actions at serotonin receptors. *J Neurosci.* 2009; 29:7124–7136. [PubMed: 19494135]
36. Alexander GM, Rogan SC, Abbas AI, Armbruster BN, Pei Y, Allen JA, Nonneman RJ, Hartmann J, Moy SS, Nicolelis MA, McNamara JO, Roth BL. Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. *Neuron.* 2009; 63:27–39. [PubMed: 19607790]
37. Cohen MS, Zhang C, Shokat KM, Taunton J. Structural bioinformatics-based design of selective, irreversible kinase inhibitors. *Science.* 2005; 308:1318–1321. [PubMed: 15919995]
38. Kenakin, T. *A Pharmacology Primer: Theory, Application, and Methods.* Third ed. Elsevier Science & Technology Books; 2009.
39. Schiffer HH, Reding EC, Fuhs SR, Lu Q, Piu F, Wong S, Littler PL, Weiner DM, Keefe W, Tan PK, Nash NR, Knapp AE, Olsson R, Brann MR. Pharmacology and signaling properties of epidermal growth factor receptor isoforms studied by bioluminescence resonance energy transfer. *Mol Pharmacol.* 2007; 71:508–518. [PubMed: 16968809]
40. Tallquist M, Kazlauskas A. PDGF signaling in cells and mice. *Cytokine Growth Factor Rev.* 2004; 15:205–213. [PubMed: 15207812]
41. Mackeigan JP, Murphy LO, Dimitri CA, Blenis J. Graded mitogen-activated protein kinase activity precedes switch-like c-Fos induction in mammalian cells. *Mol Cell Biol.* 2005; 25:4676–4682. [PubMed: 15899869]
42. Garnovskaya MN, Nebigil CG, Arthur JM, Spurney RF, Raymond JR. 5-Hydroxytryptamine_{2A} receptors expressed in rat renal mesangial cells inhibit cyclic AMP accumulation. *Mol Pharmacol.* 1995; 48:230–237. [PubMed: 7651356]
43. Ross R. Platelet-derived growth factor. *Lancet.* 1989; 1:1179–1182. [PubMed: 2566744]
44. De Meyts P, Whittaker J. Structural biology of insulin and IGF1 receptors: implications for drug design. *Nat Rev Drug Discov.* 2002; 1:769–783. [PubMed: 12360255]
45. Wong RW, Guillaud L. The role of epidermal growth factor and its receptors in mammalian CNS. *Cytokine Growth Factor Rev.* 2004; 15:147–156. [PubMed: 15110798]
46. Cornea-Hebert V, Riad M, Wu C, Singh SK, Descarries L. Cellular and subcellular distribution of the serotonin 5-HT_{2A} receptor in the central nervous system of adult rat. *J Comp Neurol.* 1999; 409:187–209. [PubMed: 10379914]
47. Mei L, Xiong WC. Neuregulin 1 in neural development, synaptic plasticity and schizophrenia. *Nat Rev Neurosci.* 2008; 9:437–452. [PubMed: 18478032]
48. Willins DL, Deutch AY, Roth BL. Serotonin 5-HT_{2A} receptors are expressed on pyramidal cells and interneurons in the rat cortex. *Synapse.* 1997; 27:79–82. [PubMed: 9268067]
49. Huang YZ, Won S, Ali DW, Wang Q, Tanowitz M, Du QS, Pelkey KA, Yang DJ, Xiong WC, Salter MW, Mei L. Regulation of neuregulin signaling by PSD-95 interacting with ErbB4 at CNS synapses. *Neuron.* 2000; 26:443–455. [PubMed: 10839362]
50. Buckley PF, Mahadik S, Pillai A, Terry A Jr. Neurotrophins and schizophrenia. *Schizophr Res.* 2007; 94:1–11. [PubMed: 17524622]
51. Kalkman HO. Altered growth factor signaling pathways as the basis of aberrant stem cell maturation in schizophrenia. *Pharmacol Ther.* 2009; 121:115–122. [PubMed: 19046988]

52. Roth BL, Willins DL, Kroeze WK. G protein-coupled receptor (GPCR) trafficking in the central nervous system: relevance for drugs of abuse. *Drug Alcohol Depend.* 1998; 51:73–85. [PubMed: 9716931]
53. Gavi S, Shumay E, Wang HY, Malbon CC. G-protein-coupled receptors and tyrosine kinases: crossroads in cell signaling and regulation. *Trends Endocrinol Metab.* 2006; 17:48–54. [PubMed: 16460957]

**Figure 1.**

RSK2 is required for inhibitory cross-talk between the EGFR and the 5-HT_{2A} receptor. The EGFR and RSK2 were activated with 100ng/mL EGF (A) and then 5-HT_{2A}-mediated Ca^{2+} responses were measured via FLIPR^{tetra} assays in RSK2^{+/+} (B-D) and RSK2^{-/-} (E-F) MEFs. A, Immunoblots showing that EGF treatment (100ng/mL) activated the EGFR (Tyr(P)-1068, top panels) and RSK2 (Ser(P)-386, bottom panels) in RSK2^{+/+} MEFs. The EGFR was similarly activated in RSK2^{-/-} MEFs; however, RSK2 was not detected in RSK2^{-/-} MEFs (bottom panel). Shown are representative immunoblots of three independent experiments. B, In RSK2^{+/+} MEFs, CRCs for 5-HT were significantly shifted rightward (i.e., decreased 5-HT potency, bold arrow) following 30 min (○), and 60 min (▲) EGF treatments relative to untreated cells (■). Shown are the results (mean \pm SEM) of

three to ten independent experiments performed in duplicate ($p < 0.05$). *C*, In RSK2^{+/+} MEFs, 5-HT_{2A}-mediated Ca²⁺ responses elicited by an EC₅₀ concentration of 5-HT (10 nM) were significantly attenuated following 30 min (○) and 60 min (▲) EGF treatments relative to untreated cells (■). Shown are normalized Ca²⁺ traces (untreated set to 100%, mean ± SEM) of three to nine independent experiments ($p < 0.05$). *D*, In RSK2^{+/+} MEFs, EGF attenuated 5-HT_{2A}-mediated Ca²⁺ responses with an IC₅₀ of 1.3 ng/mL. Shown are the results (mean ± SEM) of four independent experiments performed in duplicate. *E*, In RSK2^{-/-} MEFs, CRCs for 5-HT were not significantly shifted following EGF treatments ((30 min (○) and 60 min (▲)) relative to untreated cells (■). Shown are the results (mean ± SEM) of six independent experiments performed in duplicate ($p > 0.05$). *F*, In RSK2^{-/-} MEFs, 5-HT_{2A}-mediated Ca²⁺ responses elicited by an EC₅₀ concentration of 5-HT (10 nM) were not decreased after 30 min (○) and 60 min (▲) EGF treatments relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean ± SEM) of three to six independent experiments ($p > 0.05$).

**Figure 2.**

Genes involved in EGFR signal transduction are expressed at similarly in RSK2^{+/+} and RSK2^{-/-} MEFs. The microarray data quantifying gene expression in RSK2^{+/+} and RSK2^{-/-} MEFs was published previously by Sheffler et al. (13). Here we overlaid the mRNA expression levels of EGFR signal transduction genes in RSK2^{+/+} and RSK2^{-/-} MEFs with gene-expression color criterion and fold-changes from the programs GenMAPP and MAPPFinder. Gray colored genes are equally expressed in RSK2^{+/+} and RSK2^{-/-} MEFs. Green colored genes show greater than a 2-fold increase in expression in RSK2^{-/-} MEFs compared to RSK2^{+/+} fibroblasts. Red colored genes show greater than a 2-fold decrease in expression in RSK2^{-/-} MEFs compared to RSK2^{+/+} fibroblasts.




Figure 3.

RSK2 is required for TGF- α -induced attenuation of 5-HT_{2A} receptor signaling. The EGFR was activated with 100ng/mL TGF- α and then 5-HT_{2A}-mediated Ca²⁺ responses were measured via FLIPR^{tetra} assays in RSK2^{+/+} (A-C) and RSK2^{-/-} (D-E) MEFs. **A**, In RSK2^{+/+} MEFs, the CRC for 5-HT was significantly shifted rightward (i.e., decreased potency, bold arrow) and downward (i.e., decreased maximal signaling, bold arrow) following 60 min (o) TGF- α treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of five independent experiments performed in duplicate ($p < 0.05$). **B**, In RSK2^{+/+} MEFs, activation of 5-HT_{2A} receptors with an EC₅₀ concentration of 5-HT (10 nM) was significantly attenuated following 60 min (o) TGF- α treatment relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean \pm SEM) of five independent experiments ($p < 0.05$). **C**, In RSK2^{+/+} MEFs, TGF- α attenuated 5-HT_{2A}-mediated Ca²⁺ responses with an IC₅₀ of 4.1ng/mL. Shown are the results (mean \pm SEM) of three independent experiments performed in duplicate. **D**, In RSK2^{-/-} MEFs, the CRC for 5-HT was not significantly shifted following 60 min (o) TGF- α treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of three independent experiments performed in duplicate ($p > 0.05$). **E**, In RSK2^{-/-} MEFs, activation of 5-HT_{2A} receptors with an EC₅₀ concentration of 5-HT (10 nM) was not significantly decreased following 60 min (o) TGF- α treatment relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean \pm SEM) of three independent experiments ($p > 0.05$).




Figure 4.

The unique pharmacology of partial agonists shows that RSK2 is required for EGFR-mediated attenuation of 5-HT_{2A} receptor signaling. The EGFR was activated with 100ng/mL EGF and then 5-HT_{2A}-mediated Ca²⁺ responses elicited by the weak partial agonist lisuride (A-B) and the strong partial agonist 5-methoxyDMT (C-D) were measured via FLIPR^{tetra} assays in RSK2^{+/+} and RSK2^{-/-} MEFs. **A**, In RSK2^{+/+} MEFs, the CRC for lisuride was significantly shifted downward (i.e., decreased maximal signaling, bold arrow) following 60 min (o) EGF treatment relative to untreated cells (■). Shown are the results (mean +/- SEM) of three independent experiments performed in duplicate ($p < 0.05$). **B**, In RSK2^{-/-} MEFs, the CRC for lisuride was not significantly decreased following EGF treatment relative to untreated cells (■). Shown are the results (mean +/- SEM) of three independent experiments performed in duplicate ($p > 0.05$). **C**, In RSK2^{+/+} MEFs, the CRC for 5-methoxyDMT was significantly shifted rightward (i.e., decreased potency, bold arrow) following 60 min (o) EGF treatment relative to untreated cells (■). Shown are the results (mean +/- SEM) of three independent experiments performed in duplicate ($p < 0.05$). **D**, In RSK2^{-/-} MEFs, the CRC for 5-methoxyDMT was not significantly shifted following EGF treatment relative to untreated cells (■). Shown are the results (mean +/- SEM) of three independent experiments performed in duplicate ($p > 0.05$).





Figure 5.

RSK2 is required for inhibitory cross-talk between the PDGFR and the 5-HT_{2A} receptor. 5-HT_{2A}-mediated Ca²⁺ responses were measured via FLIPR^{tetra} assays (A) in RSK2^{+/+} and RSK2^{-/-} mVSMCs following treatment with the PDGF ligands PDGF AB (B-C) and PDGF BB (D-G). A, 5-HT elicits robust Ca²⁺ responses in RSK2^{+/+} mVSMCs which can be blocked with the 5-HT_{2A}-specific antagonist MDL100907 (100nM). Shown are the results (mean \pm SEM) of two independent experiments performed in duplicate. B, In RSK2^{+/+} mVSMCs, the CRC for 5-HT was significantly shifted downward (i.e., decreased maximal signaling, bold arrow) following 60 min (o) PDGF AB treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of four to six independent experiments performed in duplicate ($p < 0.05$). C, In RSK2^{-/-} mVSMCs, the CRC for 5-HT was not significantly shifted following PDGF AB treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of four independent experiments performed in duplicate ($p > 0.05$). D, In RSK2^{+/+} mVSMCs, the CRC for 5-HT was significantly shifted downward (i.e., decreased maximal signaling, bold arrow) following 60 min (o) PDGF BB treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of four to six independent experiments performed in duplicate ($p < 0.05$). E, In RSK2^{+/+} mVSMCs, 5-HT_{2A}-mediated Ca²⁺ responses elicited by a maximal concentration of 5-HT (10 μ M) were significantly attenuated following 60 min (o) PDGF BB treatment relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean \pm SEM) of four independent experiments ($p < 0.05$). F, In RSK2^{-/-} mVSMCs, the CRC for 5-HT was not significantly shifted following PDGF BB treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of four independent experiments performed in duplicate ($p > 0.05$). G, In RSK2^{-/-} mVSMCs, 5-HT_{2A}-mediated Ca²⁺ responses elicited by a maximal concentration of 5-HT (10 μ M) were not significantly decreased following PDGF BB treatment relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean \pm SEM) of four independent experiments ($p > 0.05$).

**Figure 6.**

IGF-1 weakly activates RSK2 and does not attenuate 5-HT_{2A} receptor signaling. 5-HT_{2A}-mediated Ca²⁺ responses were measured via FLIPR^{tetra} assays in RSK2^{+/+} and RSK2^{-/-} MEFs following treatment with 10 nM IGF-1 (A and B) or 100 ng/mL EGF (C). For Western blotting, a phospho-specific antibody was used to detect activation of RSK2 (D and E). A and B, In RSK2^{+/+} and RSK2^{-/-} MEFs, CRCs for 5-HT were not significantly shifted following 30 min (○) and 60 min (▲) IGF-1 treatments relative to untreated cells (■). Shown are the results (mean ± SEM) of three independent experiments performed in duplicate ($p > 0.05$). C, Control experiments in RSK2^{+/+} MEFs showed that 5-HT_{2A}-mediated Ca²⁺ responses elicited by an EC₅₀ concentration of 5-HT (10 nM) were significantly attenuated following 60 min (○) 100 ng/mL EGF treatment relative to untreated cells (■). Shown are the normalized Ca²⁺ traces (untreated set to 100%, mean ± SEM) of three independent experiments performed in duplicate ($p < 0.05$). D, Immunoblot showing that EGF robustly activated RSK2 (Ser(P)-386) in RSK2^{+/+} MEFs; whereas IGF-1 weakly activated RSK2 in RSK2^{+/+} MEFs. Shown are representative data from three independent experiments. E, Quantification of immunoblots in (D) showing that maximal activation of RSK2 at 5 min was significantly greater after EGF treatment than after IGF-1 treatment (*, $p < 0.05$). Shown are the results (sum pixel intensity normalized to total RSK2, mean ± SEM) of three independent experiments.




Figure 7.

RSK2 is required for inhibitory cross-talk between the EGFR and the P2Y purinergic receptor. The EGFR was activated with 100ng/mL EGF and then P2Y-mediated Ca^{2+} responses were measured via FLIPR^{tetra} assays in RSK2^{+/+} (A) and RSK2^{-/-} (B) MEFs. A, In RSK2^{+/+} MEFs, the CRC for ATP was significantly shifted rightward (i.e., decreased ATP potency, bold arrow) and downward (i.e., decreased maximal signaling, bold arrow) following 60 min (○) EGF treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of five independent experiments performed in duplicate ($p < 0.05$). B, In RSK2^{-/-} MEFs, the CRC for ATP was not significantly shifted following EGF treatment relative to untreated cells (■). Shown are the results (mean \pm SEM) of five independent experiments performed in duplicate ($p > 0.05$).




Figure 8.

Growth factors essential for normal brain function attenuate 5-HT_{2A} receptor signaling in cortical neurons. Cortical neurons were isolated from RSK2^{+/+} mice and cultured for ~14 days *in vitro* (scale bar is 20μm). These neurons were loaded with Ca²⁺ dye for 60 min prior to stimulation with the 5-HT_{2A}-selective agonist DOI (1μM) and 5-HT_{2A}-mediated Ca²⁺ responses (see color scale on right) were measured by live cell fluorescence microscopy in uninfected cortical neurons (A-C) and cortical neurons infected with GFP-tagged 5-HT_{2A} receptors (D-I). Growth factor-mediated changes in signaling were detected by comparing Ca²⁺ responses in untreated neurons (D-E) and neurons treated with 100ng/mL EGF (G-I) and 100ng/mL NRG-1 (images not shown). Ca²⁺ responses were quantified by segmenting the total area of responding cells and assessing the mean fluorescence intensity/area (F, I, J). *A and B*, The 5-HT_{2A}-selective agonist DOI did not produce measurable Ca²⁺ responses in uninfected neurons. *C*, The uninfected neurons produced measurable Ca²⁺ responses upon depolarization with 80mM KCl. *D*, Shown are cortical neurons that were infected with

lentivirus encoding GFP-tagged 5-HT_{2A} receptors prior to stimulation with DOI. *E and F*, DOI produced robust Ca²⁺ responses in infected neurons which were quantified via manual segmenting. *G*, Shown are cortical neurons infected with GFP-tagged 5-HT_{2A} receptors loaded with Ca²⁺ imaging dye containing 100ng/mL EGF for 60 min prior to stimulation with DOI. *H and I*, The 5-HT_{2A}-selective agonist DOI produced weak Ca²⁺ responses in infected neurons treated with EGF. *J*, The DOI-induced 5-HT_{2A} Ca²⁺ responses in infected neurons (○) was significantly greater than those measured from neurons pretreated with EGF (▲) or NRG-1 (▼) ($p < 0.05$). No response was detected for uninfected neurons (■). Quantified results (mean \pm SEM) are shown from three independent experiments.

Effects of various growth factors on the signaling of GPCR ligands in MEFs.

 e_{ND} , not determined.

Table 2

Effects of PDGFR agonists on 5-HT_{2A} receptor signaling in mVSMCs.

Cell type	Time treated with growth factor	RSK2 ^{+/+} mVSMCs				RSK2 ^{-/-} mVSMCs			
		0 min	30 min	60 min		0 min	30 min	60 min	
RTK ligand	GPCR ligand	$E_{max} \pm SEM, \%c$	$pEC_{50} \pm SEM^{a,b}$	$E_{max} \pm SEM, \%c$		$pEC_{50} \pm SEM^{a,b}$	$pEC_{50} \pm SEM^{a,b}$	$pEC_{50} \pm SEM^{a,b}$	$E_{max} \pm SEM, \%c$
PDGF AB	5-HT	6.91 +/- 0.05	99.4 +/- 2.2	90.5 +/- 5.4	83.9 +/- 5.6 ^d	6.80 +/- 0.10	6.38 +/- 0.09	6.66 +/- 0.22	101 +/- 10.8
PDGF BB	5-HT	7.00 +/- 0.06	99.0 +/- 2.7	94.5 +/- 7.3	79.2 +/- 5.1 ^d	6.70 +/- 0.08	6.57 +/- 0.14	6.64 +/- 0.19	95.8 +/- 7.7

^a Agonist-induced Ca²⁺ responses were quantified via FLIPR assays in untreated and growth factor-treated RSK2^{+/+} and RSK2^{-/-} mVSMCs. The fit parameters of potency (EC₅₀) and maximal signaling (E_{max}) were obtained from nonlinear regression (GraphPad software) and represent the mean +/- SEM of at least three independent experiments performed in duplicate.

^b pEC₅₀ values are represented as -log of EC₅₀ in M.

^c The maximum response of agonist (E_{max}) in untreated cells was set equal to 100%.

^d The F test was used to determine the statistical significance (defined as p<0.05) of the fit parameters in growth factor-treated and untreated cells.

Table 3Effects of EGFR family agonists on 5-HT_{2A} receptor signaling in cortical neurons.

RTK ligand	GPCR ligand	Peak Ca ²⁺ release ^a	Area under curve
None	DOI	0.67 +/- 0.03	33.7 +/- 1.3
EGF	DOI	0.45 +/- 0.03 ^b	20.9 +/- 1.2 ^b
NRG-1	DOI	0.56 +/- 0.06	28.6 +/- 1.7 ^b

^a Agonist-induced Ca²⁺ responses in untreated and growth factor-treated cortical neurons were quantified using the FLIPR Ca²⁺ assay kit and BD Pathway 855 high content imaging microscope. To control for subtle differences in 5-HT_{2A}-GFP receptor expression, Ca²⁺ responses were normalized to GFP intensity/well using custom written macros for Excel (Microsoft) and Image J (U. S. National Institutes of Health). Values were expressed as fold over baseline and represent the mean +/- SEM of three independent experiments.

^b A two-tailed, paired t test was used to determine the statistical significance (defined as $p < 0.05$) of responses in growth factor-treated vs. untreated cells.